



BY APPOINTMENT TO
HER MAJESTY QUEEN ELIZABETH II
HORSE FOOD MANUFACTURERS
DODSON & HORRELL LIMITED, NANTWICH

DODSON & HORRELL

ANIMAL HEALTH, NUTRITION AND WELL-BEING

FEEDING SUCCESS

BREEDING

EQUILAC MARES MILK



Equilac is a complementary milk replacer for rearing or weak foals

- Dissolves completely
- Optimum nutritional support
- Supporting growth and development of foals
- Can be used as an addition to mares milk
- Can be given from day one to weaning



Bag size: **10KG**
Shelf life: **18 Months from date of manufacture**
(store below 25 degrees (c))

Product Profile

Profile	
Protein	26%
Oil	12%
Ash	7%
Fibre	0%
Sodium	0.5%
Calcium	0.9%
Phosphorus	0.7%
Lactose	47%

Vitamins	
Vitamin A (3a672a)	25,000iu/kg
Vitamin D3 (3a671)	7,850iu/kg
Vitamin E (3a700)	146mg/kg
Vitamin K3 (3a711)	2mg/kg
Vitamin B1 (3a821)	2.8mg/kg
Vitamin B2 Riboflavine	8mg/kg
Nicotinic Acid (3a315)	21mg/kg
Pantothenic acid	22.5mg/kg
Vitamin B6 (3a831)	2.9mg/kg
Vitamin B12	29.3mcg/kg
Biotin (3a880)	100mcg/kg
Folic Acid (3a316)	0.46mg/kg
Vitamin C (3a300)	50mg/kg

Mineral values for Equilac are total values within the product including added and naturally occurring minerals from grains, fibres and other raw materials. Such materials are naturally variable and as such total values may vary slightly.

Statutory Information

Additives: Nutritional	
Iron (Iron (II) Sulphate Monohydrate/3b103)	53mg/kg
Iodine (Potassium Iodide/3b201)	2.5mg/kg
Copper (Copper (II) sulphate pentahydrate/3b405)	5mg/kg
Manganese (Manganous sulphate monohydrate/3b503)	40mg/kg
Zinc (Zinc sulphate monohydrate/3b605)	40mg/kg
Selenium (Sodium selenite/3b801)	0.25mg/kg

Equilac Composition

Skim milk powder, fat filled skim milk powder, whey powder, fat filled whey powder, dextrose.

Uses

Equilac Mare's Milk Replacer can be used for hand rearing, supplementary feeding or automatic machine feeding.

Hand Rearing

If a foal is orphaned or the mare has no milk, Equilac mares milk Replacer is suitable to feed from one day old up to weaning. Equilac provides calories, protein, vitamins and minerals to support the growing foal.

Equilac is not a source of colostrum. It is vital that colostrum is consumed as the primary source of milk within the first 12-14 hours of birth.

Supplementary Feeding

When a mare has insufficient milk, which can frequently happen at the start of lactation, Equilac Milk Replacer should be fed as a milk supplement.

Feeding Rate

Equilac mixes instantly in warm water (130g per litre) – add the milk powder to the water and whisk. Equilac should be fed at 37-38 °C do not reheat.

For complete accuracy weigh Equilac before mixing. These instructions should be used as a guide only, adjusting the quantity fed according to bodyweight and condition.

For further detailed feeding instructions please contact us via helplineenquiries@dodsonandhorrell.com

For equine use only. A small amount of blue specs in Equilac should be expected and is perfectly normal, due to the inclusion of a bioavailable source of copper.

For more help, advice or information on nutrition for your horse or finding the right feed for your horse, please contact us.

Nutritional helpline: 01270 782223
enquiries@dodsonandhorrell.com
dodsonandhorrell.com