Ingredients

Dextrose, Citric Acid, Ascorbic Acid (Vitamin C), Magnesium Sulfate, D-Calcium Pantothenate, Potassium Chloride, Sodium Chloride, Iron Sulfate, Silicon Dioxide, Folic Acid, Vitamin A Acetate, D-Activated Animal Sterol (Vitamin D3), Vitamin E Supplement, Menadione Sodium Bisulfite Complex (source of Vitamin K), Niacinamide, Riboflavin Supplement, Pyridoxine Hydrochloride, Thiamine Hydrochloride, Vitamin B12 Supplement, Dried Lactobacillus acidophilus Fermentation Product, and Dried Enterococcus faecium Fermentation Product.

Keep out of reach of children. For Animal Use Only.

STORE IN A COOL, DRY PLACE

®Registered Trademark of Agri Laboratories, Ltd.



Manufactured for: Agri Laboratories, Ltd. St. Joseph, MO 64503

Vitamins & Electrolytes "Plus"

Water Soluble Powder with Vitamins, Electrolytes, and Acidifiers for Livestock and Poultry

Guaranteed Analysis

(minimum, unless otherwise stated)

| | Per 4 oz | Per Ib |
|---------------------------|--------------|---------------|
| Salt | 11.0% (min) | 11.0% (min) |
| | 13.2% (max) | 13.2% (max) |
| Sodium | 4.70% | 4.70% |
| Potassium | 3.80% | 3.80% |
| Magnesium | 0.26% | 0.26% |
| Vitamin A | 2,500,000 IU | 10,000,000 IU |
| Vitamin D₃ | 1,000,000 IU | 4,000,000 IU |
| Vitamin E | 2,000 IU | 8,000 IU |
| Riboflavin | 750 mg | 3,000 mg |
| Pantothenic Acid | 2,000 mg | 8,000 mg |
| Folic Acid | 125 mg | 500 mg |
| Thiamine | 375 mg | 1,500 mg |
| Niacin | 5,000 mg | 20,000 mg |
| Pyridoxine | 300 mg | 1,200 mg |
| Ascorbic Acid (Vitamin C) | 3,750 mg | 15,000 mg |
| Vitamin B ₁₂ | 2.5 mg | 10 mg |
| Menadione (Vitamin K) | 1,000 mg | 4,000 mg |

NET CONTENTS: 4 oz (112 gm) PACK





Suggested Use

Use on Cattle, Swine, Sheep, Horses, Poultry and Turkeys during periods of stress or reduced feed intake

Directions for Use

Drinking Water: Mix contents of one pouch (4 oz) in 128 gallons of water.

For Automatic Proportioners:

Mix contents of one pouch (4 oz) in 1 gallon of water to prepare a stock solution. Set the proportioner to deliver 1 fl oz of stock solution per 1 gallon of drinking water. Prepare fresh stock solution daily.

Feed Use: Mix contents of one pouch (4 oz) per 500 lb of complete feed.

Provide for 3-5 days, repeat as needed.

